

Parasites of the Southern silvery grebe *Podiceps occipitalis* (Aves, Podicipedidae) in Chile

Parasitas do mergulhão-de-orelha-amarela *Podiceps occipitalis* (Aves, Podicipedidae) no Chile

Daniel González-Acuña¹; Sebastián Llanos-Soto²; Carlos Landaeta-Aqueveque¹; Felipe González¹;
John Mike Kinsella³; Sergey Mironov⁴; Armando Cicchino⁵; Carlos Barrientos⁶; Gonzalo Torres-Fuentes¹;
Lucila Moreno^{2*}

¹ Facultad de Ciencias Veterinarias, Universidad de Concepción, Chillán, Chile

² Facultad de Ciencias Naturales y Oceanográficas, Universidad de Concepción, Concepción, Chile

³ Helm West Lab, Missoula, MT, USA

⁴ Zoological Institute, Russian Academy of Sciences, Universitetskaya Embankment 1, Saint Petersburg, Russia

⁵ Universidad Nacional de Mar del Plata, Mar del Plata, Argentina

⁶ Escuela de Medicina Veterinaria, Universidad Santo Tomás, Concepción, Chile

Received January 17, 2017

Accepted March 6, 2017

Abstract

A total of 97 southern silvery grebes (*Podiceps occipitalis*), which died as the result of an oil spill on the coast of central Chile, were examined for ecto- and endoparasites. Two lice species including *Aquanirmus rollandii* (Phloptoridae) and *Pseudomenopon dolium* (Menoponidae) were found from 6.2% (6/97) of birds. In 91.7% (89/97) of cases, grebes were infected with some kind of helminths. Three species of gastrointestinal helminths were detected: *Eucoelus contortus* (Nematoda), *Profilicollis bullocki* (Acanthocephala), and *Confluaria* sp. (Cestoda). In addition, *Pelecitus fulicaeatrae* (Nematoda) was removed from the tibiotarsal-tarsometatarsal articulation in 13.4% (13/97) of the specimens examined. To our knowledge, these are the first records of *A. rollandii*, *E. contortus*, and *Confluaria* sp. as parasites of *P. occipitalis*. In addition, these findings expand the distributional range of *A. rollandii*, *E. contortus*, *P. fulicaeatrae*, and *Confluaria* sp. to Chile.

Keywords: Acanthocephalans, cestode, nematode, ectoparasites, endoparasites, water birds.

Resumo

Um total de 97 mergulhões-de-orelha-amarela (*Podiceps occipitalis*), que morreram devido a um derramamento de óleo na costa do Chile central, foram examinados em busca de parasitos internos e externos. Parasitos externos foram encontrados em 6,2% (6/97) das aves, com a identificação de duas espécies de piolhos: *Aquanirmus rollandii* (Phloptoridae) e *Pseudomenopon dolium* (Menoponidae). Em 91,7% (89/97) dos casos, os mergulhões apresentaram algum tipo de parasito interno. Foram detectadas três espécies de parasitos gastrointestinais: *Eucoelus contortus* (Nematoda), *Profilicollis bullocki* (Acanthocephala) e *Confluaria* sp. (Cestoda). Além disso, *Pelecitus fulicaeatrae* (Nematoda) foi isolado das articulações tibiotársica e tarsometatarsal em 13,4% (13/97) das aves examinados. Estes resultados correspondem ao primeiro relato de *A. rollandii*, *E. contortus* e *Confluaria* sp. associados com *P. occipitalis*, e expandem a distribuição destes parasitos e *P. fulicaeatrae* para o Chile.

Palavras-chave: Acantocéfala, cestoda, nematoides, parasitos externo, parasitos interno, aves aquáticas.

*Corresponding author: Lucila Moreno Salas. Facultad de Ciencias Naturales y Oceanográficas, Universidad de Concepción, Casilla 160C, Concepción, Chile. e-mail: lumoreno@udec.cl.

Introduction

The southern silvery grebe *Podiceps occipitalis* Garnot, 1826 is widely distributed across South American countries, as it can be found in Argentina (including in the Falkland Islands/Malvinas), Bolivia, Peru, Paraguay, and Chile, although they rarely occur in Ecuador and Colombia (COUVE & VIDAL, 2003; MAILLARD et al., 2006). In Chile, grebes are represented by two subspecies, *P. occipitalis occipitalis*, which distributed from Atacama to Tierra del Fuego, and *P. occipitalis juninensis*, which inhabits lakes at higher altitudes (3500-4500 m) in the Andean plateau (JARAMILLO, 2005). In terms of its conservation status, this species has been categorized as of least concern (LC); however, the overall population appears to be decreasing overall (BIRDLIFE INTERNATIONAL, 2016). Furthermore, *P. occipitalis* is vulnerable to oil spill events due to their gregarious behavior during the winter, as they gather in large water bodies, sea bays, and lagoons to form flocks consisting of hundreds of individuals (JARAMILLO, 2005). Knowledge about parasite diversity and prevalence in *P. occipitalis* is quite limited (HINOJOSA-SÁEZ & GONZÁLEZ-ACUÑA, 2005; ATKINSON et al., 2008). To date, only a louse, *Pseudomenopon dolium* Rudow, 1896 (Phthiraptera: Menopodidae), and helminths, *Pelecitus fulicaeatrae* Diesing, 1861 (Nematoda: Filarioidea) and *Proflicollis bullocki* Perry, 1942 (Acanthocephala: Polymorphidae), have been described for *P. occipitalis* in Bolivia, Argentina, and Chile, respectively (PRICE et al., 2003; RIQUELME et al., 2006; ESCUDERO et al., 2007). The present study provides new data on the diversity and prevalence of ecto- and endoparasites in the southern silvery grebe that inhabits the coast of central Chile.

Materials and Methods

In May 2007, an oil spill event occurred along the coast of central Chile, near Lenga Town (36° 46' S, 73° 10' W). In order to protect local wildlife, different seabird species were captured and transported to the Wildlife Rehabilitation Center from the Universidad de Concepción, Chillán, so they could be rehabilitated. Unfortunately, 97 grebes (*P. occipitalis occipitalis*) died while being washed with detergent in order to remove the oil from their feathers. Their carcasses were moved to the Animal Science Department at the Universidad de Concepción and each bird was kept individually frozen at -12 °C for further analysis.

The grebes were externally inspected and their feathers were closely examined for ectoparasites. Following collection, the ectoparasites were preserved in 70% ethanol. Lice were mounted using Canada balsam following the technique of Price et al. (2003); they were identified under a light microscope based on the keys and descriptions of Castro & Cicchino (2000) and Price et al. (2003). Specimens were photographed via microscope and measured using Micrometrics® (Micrometrics Instrument Corporation, Norcross, GA, USA).

For endoparasites, the carcasses were necropsied using the protocols detailed in Kinsella & Forrester (1972). Endoparasites were identified following the descriptions provided by Betlejewska et al. (2002), Escudero et al. (2007), Vasileva et al. (1999a, b, 2000, 2001), and Mateo et al. (1982).

Results and Discussion

Lice including *Aquanirmus rollandii* Castro & Cicchino, 2000 (Phthiraptera, Ischnocera) (Figure 1A) (2 females, 1 male, and 2 nymphs) and *Pseudomenopon dolium* (Figure 1B) (5 females and 2 males) were found on 6.2% (6/97) of grebes (Table 1). Both parasite species were restricted to Podicipedidae. With respect to *A. rollandii*, it was only previously reported in *Rollandia rolland chilensis* (Lesson) in Argentina (CASTRO & CICHINO, 2000). In contrast, *P. dolium* is a cosmopolitan species that parasitizes *Podiceps ruficollis*, *P. auritus*, *P. griseigena*, *P. cristatus*, *P. nigricollis*, *Podilymbus podiceps*, *Aechmophorus occidentalis*, and *Tachybaptus ruficollis* in Europe, Africa, India, Asia and North America (PRICE, 1974; MARTÍN MATEO, 2006; VAS et al., 2012; DIK & HALAJIAN, 2013; GALLOWAY et al., 2014; JAŁOSZYŃSKI et al., 2014). In South America, *P. dolium* has been described as a parasite of *P. podiceps* in Argentina, *R. rolland chilensis* in Chile and Argentina, *P. taczanowskii* in Peru, and *P. occipitalis* in Bolivia (PRICE, 1974; CICHINO, 2011).

Only a small number of individuals from both species were collected. This could be explained by the fact that grebes were covered in oil and subsequently washed with detergent to remove it. For this reason, our study may be underestimating the intensity of infection in the examined birds. Nonetheless, this marks the first time that *A. rollandii* has been identified from *P. occipitalis*, and where its distributional range has expanded to Chile.

For the same reason, feather mites (Acari: Analgoidea and Pterolichoidea), the most abundant and diverse arthropods living on the plumage and bodies of birds, were not detected on the examined carcasses of *P. occipitalis*. Nevertheless, while taking into account the currently known data on feather mites associated with Podicipedidae, it is probable that representatives of at least three different feather mite families (Ptiloxenidae, Xolalgidae and Laminosioptidae) will be found on the southern silvery grebe in Chile. Thus, feather mites of the genus *Ptiloxenus* Hull, 1934 (Pterolichoidea: Ptiloxenidae) was associated with grebes of the genera *Podiceps* and *Rollandia* (DABERT & EHRNSBERGER, 1998). *Ingrassia colymbi* Gaud, 1974 (Analgoidea: Xolalgidae), the only




Figure 1. *Aquanirmus rollandii* female (A) and *Pseudomenopon dolium* female (B) (100× magnification).

Table 1. Summary of ecto- and endoparasites identified on the southern silvery grebe *Podiceps occipitalis* from Central Chile.

Parasite	Location	Prevalence (%)	Total parasites	Range	Mean Intensity	Mean Abundance
Endoparasites						
<i>Eucoleus contortus</i>	Small intestine	1.03	1	1	1	0.01
<i>Confluaria</i> sp.	Small intestine	46.39	708	1-141	15.7	7.30
<i>Profilicollis bullocki</i>	Small intestine	68.04	975	1-81	14.7	10.05
<i>Pelecitus fulicaeatrae</i>	Tibiotarsal-tarsometatarsal articulation	13.40	72	1-18	5.5	0.70
Ectoparasites						
<i>Aquanirmus rollandii</i>	Feathers	3.09	5	1-3	0.05	1.60
<i>Pseudomenopon dolium</i>	Feathers	3.09	7	1-3	0.07	2.30

representative of the family associated with grebes, was described from the covert feathers of the bodies of *Tachybaptus ruficollis* (GAUD, 1974). A quill-wall mite known as *Podicipedicoptes americanus* (LOMBERT et al., 1979) (Analgoidea: Laminosioptidae), the only representative of the genus *Podicipedicoptes*, was described from *Podilymbus podiceps* (LOMBERT et al., 1979). Species of the genus *Podicipedicoptes*, *Ptiloxenus* and *Ingrassia* were also expected to be found on the southern silvery grebe.

Gastrointestinal helminths were found from 91.7% of southern silvery grebe. In addition, *Pelecitus fulicaeatrae* (Figure 2) was removed from the tibiotarsal-tarsometatarsal articulation in a smaller number of birds (13.4% of cases) (Table 1). A female of *Eucoleus contortus* (syn. *Capillaria contorta*) was collected from the small intestine of a single grebe. This parasite is normally found in the upper digestive tracts of birds (BETLEJEWSKA et al., 2002) and, in some cases, in the proventriculus (MONTEIRO et al., 2011). It commonly affects domestic Galliformes, but it also parasitizes captive-raised wild birds (MILLÁN et al., 2002; CRUZ et al., 2016). For instance, *E. contortus* has been described in swans, ducks, gulls, bustards, and cormorants in Asia, Europe and South America (Brazil) (THRELFALL, 1982; FEDYNICH et al., 1997; BOSCH et al., 2000; BETLEJEWSKA et al., 2002; VILLANÚA et al., 2007; YOSHINO et al., 2009; MONTEIRO et al., 2011; SANTORO et al., 2011). This study is the first report of *E. contortus* from grebes (Podicipedidae), and it expands the parasite's distributional range to Chile. *E. contortus* is a generalist parasite and it has been reported to parasitize other seabirds. For this reason, it is likely that *P. occipitalis* plays a role as a natural host for *E. contortus*. However, it could also represent an accidental infectious event, as only a single *E. contortus* individual was collected from the 97 grebes examined. The feeding behavior of these birds (mostly insects and aquatic crustaceans) (ROTTMANN, 1995) strengthens the hypothesis for an accidental finding of *E. contortus*, which probably was ingested by this grebe within an invertebrate host. Additionally, one must consider that *P. occipitalis* coexists with different seabirds along the coast of central Chile, including birds that were previously indicated as hosts for *E. contortus* (e.g. *Phalacrocorax brasilianus*) (MONTEIRO et al., 2011). Furthermore, this parasite was identified as a possible threat for the conservation of wild species (VILLANÚA et al., 2007), mostly due to its negative effects on the body's overall condition (BOSCH et al., 2000) and its susceptibility to predation (MILLÁN et al., 2002). For these reasons, further studies assessing parasite prevalence and diversity

**Figure 2.** *Pelecitus fulicaeatrae* removed from tibiotarsal-tarsometatarsal articulation.

should be performed in Podicipedidae to adequately elucidate their parasite fauna.

Confluaria sp. Ablasov, 1953 (Cestoda: Hymenolepididae) were collected from the small and large intestines of grebes; the latter location is unusual for these parasites, and it may be the result of parasite movements following death. Vasileva et al. (1999a) indicate that the genus *Confluaria* is restricted to Podicipedidae, and that records in other avian families are erroneous. This genus has been reported in *P. ruficollis japonicus*, *P. nigricollis*, *Tachybaptus ruficollis*, *P. grisegena*, *P. cristatus* and *P. auritus* in Asia and Europe (VASILEVA et al., 1999a, b; BARUŠ et al., 2000; VASILEVA et al., 2000, 2001, 2008; HAUKISALMI, 2015; SITKO & HENEBERG, 2015). In South America, the genus *Confluaria* is only known to be from Northern Brazil, with *P. dominicus* serving as its host (VASILEVA et al., 1999a). Therefore, this is the first time that the *Confluaria* sp. is reported from *P. occipitalis*, thus extending its range to Chile.

In this study, specimens of *Profilicollis bullocki*, which were aligned with the characteristics of this parasite found in seabirds inhabiting the coast of Central and Northern Chile (OLIVA et al., 1992; RIQUELME et al., 2006), were reported to be highly prevalent in *P. occipitalis* (68.0%). Indeed, Riquelme et al. (2006) suggest that *P. occipitalis* may act as a natural reservoir for *P. bullocki* during the winter, thus playing a role in the maintenance of infections in seabird communities during that season. This parasite was previously

identified in various charadriiform birds, including *Calidris* sp., *Larus modestus*, *L. serranus*, *L. dominicanus*, *L. pipixcan*, and *Numenius phaeopus*, as well as in *P. occipitalis* on the Pacific Coast of South America (OLIVA et al., 1992; RIQUELME et al., 2006).

In addition, *Pelecitus fulicaeatrae* was detected in 13 grebes, which were located in the tibiotarsal-tarsometatarsal articulation. Chewing lice such as *P. dolium* act as intermediate hosts (BARTLETT & ANDERSON, 1987). In the definitive host, microfilariae inhabit the feathered skin of their hosts, and adults occur in the nodules in their legs (BARTLETT & ANDERSON, 1989). Different bird families have been reported as hosts of *P. fulicaeatrae*, including grebes (VANDERBURGH et al., 1984; BARTLETT & GREINER, 1986; BARTLETT & ANDERSON, 1989; ESCUDERO et al., 2007). In South America, the occurrence of *P. fulicaeatrae* was previously reported in the gray-necked wood-rail *Aramides cajanea* (Rallidae) and in the jabiru *Jabiru myzocera* (Ciconiidae) in Brazil (PINTO & NORONHA, 2003). Escudero et al. (2007) reported the first record of *P. occipitalis* from Argentina. Our findings represent a new record of this parasite in Chile.

Our study contributes to the current knowledge on the parasite prevalence and diversity in *P. occipitalis*, which includes the first records of *A. rollandii*, *E. contortus*, and *Confluaria* sp. for this host. In addition, it expands the distributional range for *A. rollandii*, *E. contortus*, *P. fulicaeatrae*, and *Confluaria* sp. to Chile.

Acknowledgements

We appreciate the support given by the National Forest Corporation, CONAF, as well as by the Agricultural and Livestock Service, SAG, of Chile. We thank Karen Ardiles, Carolina Silva, Pedro Álvarez, Sebastián Muñoz, Braulio Muñoz, Danny Fuentes, Félix Varas, Ignacia Najle, Iván Torres, Nicolás Fernández, Pablo Olmedo, and Diego Barrientos for their support in the field. This research was funded by FONDECYT Project 1130948.

References

Atkinson CT, Thomas NJ, Hunter DB. *Parasitic diseases of wild birds*. Ames: Willey-Blackwell; 2008. <http://dx.doi.org/10.1002/9780813804620>.

Bartlett CM, Anderson RC. Mallophagan vectors and the avian filarioids: new subspecies of *Pelecitus fulicaeatrae* (Nematoda: Filarioidea) in sympatric North American hosts, with development, epizootiology, and pathogenesis of the parasite in *Fulica americana* (Aves). *Can J Zool* 1989; 67(11): 2821-2833. <http://dx.doi.org/10.1139/z89-398>.

Bartlett CM, Anderson RC. *Pelecitus fulicaeatrae* (Nematoda: Filarioidea) of coots (Gruiformes) and grebes (Podicipediformes): skin-inhabiting microfilariae and development in Mallophaga. *Can J Zool* 1987; 65(11): 2803-2812. <http://dx.doi.org/10.1139/z87-423>.

Bartlett CM, Greiner EC. A revision of *Pelecitus* Railliet & Henry, 1910 (Filarioidea, Dirofiliariinae) and evidence for the "capture" by mammals of filarioids from birds. *Bull Mus Hist Nat* 1986; 8(1): 47-99.

Baruš V, Tenora F, Kráčmar S. Heavy metal (Pb, Cd) concentrations in adult tapeworms (Cestoda) parasitizing birds (Aves). *Helminthologia* 2000; 37(3): 131-136.

Betlejewska KM, Kalisinska E, Korniyushin VV, Salamatin R. *Eucoleus contortus* (Creplin, 1839) nematode in mallard (*Anas platyrhynchos* Linnaeus, 1758) from north-western Poland. *Electron J Polish Agric Univ* 2002; 5(1): 3.

BirdLife International. *Species factsheet: Podiceps occipitalis*. Cambridge: BirdLife International; 2016 [cited 2016 June 28]. Available from: <http://www.birdlife.org/datazone/species/factsheet/62114375>

Bosch M, Torres J, Figuerola J. A helminth community in breeding Yellow-legged Gulls (*Larus cachinnans*): pattern of association and its effect on host fitness. *Can J Zool* 2000; 78(5): 777-786. <http://dx.doi.org/10.1139/cjz-78-5-777>.

Castro D, Cicchino A. A new species of *Aquanirmus* Clay and Meinertzhagen, 1939 (Phthiraptera, Philopteridae) parasitic on *Rollandia Rolland chilensis* (Lesson, 1828) (Aves, Podicipitidae), with remarks on the external chorionic morphology of the eggs. *Pap Avulsos Zool* 2000; 41(14): 213-221.

Cicchino AC. *Piojos (Insecta: Psocodea: Phthiraptera) parásitos de Gruiformes y Podicipediformes (Aves) en la Argentina* [Dissertation]. Mar del Plata: Universidad Nacional de Mar del Plata; 2011.

Couve E, Vidal C. *Birds of Patagonia, Tierra Del Fuego and Antarctic Peninsula: the Falkland Islands and South Georgia*. Punta Arenas: Fantástico Sur Birding; 2003.

Cruz CEF, Fredo G, Casagrande R, Oliveira L, Rolim V, Marques S, et al. *Eucoleus contortus* parasitism in captive-bred valley quail *Callipepla californica* (Shaw, 1798): disease and control. *Zool Gart* 2016; 85(3-4): 152-159. <http://dx.doi.org/10.1016/j.zoolgart.2016.01.008>.

Dabert J, Ehrnsberger R. Phylogeny of the feather mite family Ptiloxenidae Gaud, 1982 (Acari: Pterolichoidea). In: Ebermann E. *Arthropod biology: contributions to morphology, ecology and systematics*. 1998. p. 145-178. vol. 14. Biosystematics and Ecology Series.

Dik B, Halajian A. Chewing lice (Phthiraptera) of several species of wild birds in Iran, with new records. *J Arthropod Borne Dis* 2013; 7(1): 83-89. PMID:23785698.

Escudero G, Diaz JI, Notarnicola J. New host and distribution records of *Pelecitus fulicaeatrae* (Diesing, 1861) (Nematoda, Onchocercidae). *Acta Parasitol* 2007; 52(4): 419-421. <http://dx.doi.org/10.2478/s11686-007-0058-4>.

Fedynich AM, Pence DB, Bergan JF. Helminth community structure and pattern in sympatric populations of double-crested and Neotropic cormorants. *J Helminthol Soc Wash* 1997; 64(2): 176-182.

Galloway TD, Proctor HC, Mironov SV. Chewing Lice (Insecta: Phthiraptera: Amblycera, Ischnocera) and Feather Mites (Acari: Astigmatina: Analgoidea, Pterolichoidea): Ectosymbionts of Grassland Birds in Canada. In: Cárcamo HA, Giberson DJ. *Arthropods of Canadian Grasslands: Biodiversity and Systematics (volume 3) Part 1*. Ottawa: Biological Survey of Canada; 2014. p. 139-188.

Gaud J. Quelques espèces nouvelles de Sarcoptiformes plumicoles (Analgidae & Dermoglyphidae) parasites d'oiseaux d'Europe. *Acarologia* 1974; 15(4): 727-758. PMID:4446922.

Haukisalmi V. Checklist of tapeworms (Platyhelminthes, Cestoda) of vertebrates in Finland. *ZooKeys* 2015; 533(533): 1-61. PMID:26668540. <http://dx.doi.org/10.3897/zookeys.533.6538>.

Hinojosa-Sáez A, González-Acuña D. Estado actual del conocimiento de helmintos en aves silvestres de Chile. *Gayana (Concept)* 2005; 69(2): 241-253.

- Jałoszyński P, Gustafsson DR, Wanat A, Wanat M. Type specimens of Phthiraptera in the collection of Jadwiga Złotorzycka preserved in the Museum of Natural History, University of Wrocław. *Genus* 2014; 25(4): 645-661.
- Jaramillo A. *Aves de Chile*. Barcelona: Lynx Edicions; 2005.
- Kinsella JM, Forrester DJ. Helminths of the Florida duck, *Anas platyrhynchos fulvigula*. *Proc Helminthol Soc Wash* 1972; 39(2): 173-176.
- Lombert HAPM, Kethley JB, Lukoschus FS. Observations on quill wall mites from American birds (Acaridei: Laminosioptidae: Faincoptinae). *Int J Acarol* 1979; 5(2): 103-110. <http://dx.doi.org/10.1080/01647957908683132>.
- Maillard O, Sánchez G, Caballero E, Velásquez MÁ. Nuevo dato en la distribución de *Podiceps occipitalis* en Bolivia. *Kempfiana* 2006; 2(1): 106-108.
- Martín Mateo MP. Diversidad y distribución de las especies de Mallophaga (Insecta) en aves y mamíferos de la comunidad de Madrid. *Graellsia* 2006; 62: 21-32. <http://dx.doi.org/10.3989/graellsia.2006.v62.iExtra.108>.
- Mateo E, Córdova R, Guzmán E. *Polymorphus (Profilicollis) bullocki*, nueva especie de acantocéfalo hallado en la gaviota *Larus belcheri* en el Perú. *Bol Lima* 1982; 4(24): 73-78.
- Millán J, Gortázar C, Tizzani P, Buenestado FJ. Do helminths increase the vulnerability of released pheasants to fox predation? *J Helminthol* 2002; 76(3): 225-229. PMID:12363375. <http://dx.doi.org/10.1079/JOH2002125>.
- Monteiro CM, Amato JF, Amato SB. Helminth parasitism in the Neotropical cormorant, *Phalacrocorax brasilianus*, in southern Brazil: effect of host size, weight, sex, and maturity state. *Parasitol Res* 2011; 109(3): 849-855. PMID:21431903. <http://dx.doi.org/10.1007/s00436-011-2311-x>.
- Oliva ME, Luque JL, Cevallos A. Parásitos de *Emerita analoga* (Stimpson) (Crustacea): implicancias ecológicas. *Bol Lima* 1992; 79: 77-80.
- Pinto RM, Noronha D. Analysis of Brazilian species of *Pelecitus* Railliet & Henry (Nematoda, Filarioidea) with the establishment of new records. *Rev Bras Zool* 2003; 20(2): 361-364. <http://dx.doi.org/10.1590/S0101-81752003000200029>.
- Price RD, Hellenthal RA, Palma RL, Jonhson KP, Clayton DH. *The chewing lice: world checklist and biological overview*. Springfield: Illinois Natural History Survey; 2003.
- Price RD. A review of the genus *Pseudomenopon* (Mallophaga: Menoponidae). *Ann Entomol Soc Am* 1974; 67(1): 73-84. <http://dx.doi.org/10.1093/aesa/67.1.73>.
- Riquelme C, George-Nascimento M, Balboa L. Morfometría y fecundidad de *Profilicollis bullocki* Mateo, Córdova & Guzmán 1982 (Acantocephala: Polymorphidae) en especies simpátricas de aves costeras de Chile. *Rev Chil Hist Nat* 2006; 79(4): 465-474. <http://dx.doi.org/10.4067/S0716-078X2006000400005>.
- Rottmann J. *Guía de identificación de aves de ambientes acuáticos*. Chile: Unión de Ornitólogos de Chile; 1995.
- Santoro M, Mattiucci S, Kinsella JM, Aznar FJ, Giordano D, Castagna F, et al. Helminth community structure of the Mediterranean gull (*Ichthyaeetus melanocephalus*) in Southern Italy. *J Parasitol* 2011; 97(2): 364-366. PMID:21506790. <http://dx.doi.org/10.1645/GE-2602.1>.
- Sitko J, Heneberg P. Host specificity and seasonality of helminth component communities in central European grebes (Podicipediformes) and loons (Gaviiformes). *Parasitol Int* 2015; 64(5): 377-388. PMID:26008120. <http://dx.doi.org/10.1016/j.parint.2015.05.012>.
- Threlfall W. Endoparasites of the double-crested cormorant (*Phalacrocorax auritus*) in Florida. *Proc Helminthol Soc Wash* 1982; 49(1): 103-108.
- Vanderburgh DJ, Anderson RC, Stock TM. *Pelecitus tubercauda* n. sp. (Nematode: Filarioidea) from *Geothlypis trichas* L. and a redescription of *P. fulicaeae* (Diesing, 1861) López-Neyra, 1956. *Can J Zool* 1984; 62(3): 362-367. <http://dx.doi.org/10.1139/z84-056>.
- Vas Z, Rékási J, Rózsa L. A checklist of lice of Hungary (Insecta: Phthiraptera). *Annl Hist Nat Mus Natn Hung* 2012; 104: 5-109.
- Vasileva GP, Georgiev BB, Genov T. Palaeartic species of the genus *Confluaria* Ablasov (Cestoda, Hymenolepididae): a redescription and synonymy of *C. capillaris* (Rudolphi, 1810). *Syst Parasitol* 1999a; 43(1): 49-57. PMID:10613530. <http://dx.doi.org/10.1023/A:1006132708859>.
- Vasileva GP, Georgiev BB, Genov T. Palaeartic species of the genus *Confluaria* Ablasov (Cestoda, Hymenolepididae): redescription of *C. multistriata* (Rudolphi, 1810) and *C. japonica* (Yamaguti, 1935), and a description of *Confluaria* sp. *Syst Parasitol* 1999b; 44(2): 87-103. PMID:10619078. <http://dx.doi.org/10.1023/A:1006157504152>.
- Vasileva GP, Georgiev BB, Genov T. Palaeartic species of the genus *Confluaria* Ablasov (Cestoda, Hymenolepididae): redescription of *C. podicipina* (Szymanski, 1905) and *C. furcifera* (Krabbe, 1869), description of *C. pseudofurcifera* n. sp., a key and final comments. *Syst Parasitol* 2000; 45(2): 109-130. PMID:10743855. <http://dx.doi.org/10.1023/A:1006237509781>.
- Vasileva GP, Korniyushin VV, Genov T. Hymenolepidid cestodes from grebes (Aves, Podicipedidae) in Ukraine: the genus *Confluaria*. *Vest Zool* 2001; 35(6): 13-31.
- Vasileva GP, Skirnisson K, Georgiev BB. Cestodes of the horned grebe *Podiceps auritus* (L.) (Aves: Podicipedidae) from Lake Myvatn, Iceland, with the description of *Confluaria islandica* n. sp. (Hymenolepididae). *Syst Parasitol* 2008; 69(1): 51-58. PMID:18030602. <http://dx.doi.org/10.1007/s11230-007-9110-x>.
- Villanúa D, Casas F, Viñuela J, Gortázar C, De la Morena EG, Morales M. First occurrence of *Eucoleus contortus* in a Little Bustard *Tetrax tetrax*: negative effect of Red-legged Partridge *Alectoris rufa* releases on steppe bird conservation? *Ibis* 2007; 149(2): 405-406. <http://dx.doi.org/10.1111/j.1474-919X.2006.00620.x>.
- Yoshino T, Uemura J, Endoh E, Kaneko M, Osa Y, Asakawa M. Parasitic nematodes of anseriform birds in Hokkaido, Japan. *Helminthologia* 2009; 46(2): 117-122. <http://dx.doi.org/10.2478/s11687-009-0023-x>.